



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

1° N. α <i>Lyncis</i> .	1½° S. η <i>Leo. majoris</i> .	1½° N. β <i>Cancr.</i>
1¾° N. δ <i>Leo. min.</i>	σ “	1½° N. ζ <i>Hydræ</i> .
2¾° N. ξ <i>Virginis</i> .	β “	4½° N. μ <i>Sextan.</i>
μ “	1° S. α “	1° N. γ <i>Corvi</i> .
2¼° S. ζ <i>Ophiuchi</i> .	3½° N. <i>20 Libræ</i> .	2¼° N. γ <i>Hydræ</i> .
4½° S. ν “	Just N. π <i>Scorpii</i> .	4¾° S. π “
3½° N. <i>e Sagittarii</i> .	1° S. α “	6½° N. κ <i>Centauri</i> .
1° N. ϵ <i>Aquarii</i> .	1½° N. δ <i>Sagittarii</i> .	Just N. θ <i>Scorpii</i> .
1° N. μ “	τ “	2° N. ι <i>Sagittarii</i> .
1° N. β “	Just N. χ <i>Capricorni</i> .	1¼° S. η <i>Piscis Aust.</i>
1° S. α “	2° S. ι <i>Aquarii</i> .	ϵ “ “
4½° N. γ <i>Piscium</i> .	2½° S. λ “	Just S. <i>98 Aquarii</i> .
	R. A. 0 ^h ; Dec. 0°.	104 “

The above are the outlines as they appear to me, deduced from observations in all latitudes from 55° N. to 55° S.

It also seems to me that to a northern observer the northern boundary will appear about the same, and the southern boundary less southerly than here given, with a consequent displacement to the northward of the line of central axis of about 1° for every 19° of latitude he is distant from that part of the sky he is observing (*vice versa* in southern latitudes). At times, also, the band appears broader than here given, as if the Light extended still farther; but so faint, attenuated, and vague are such boundaries, that I have ignored them.

Extreme accuracy must not be expected, for only star-charts of a primitive nature have been employed for laying off my observations.

R. M. S. “WARRIMOO,” VANCOUVER, B. C., Dec. 4, 1899.

OBSERVATIONS OF VARIABLE STARS.

BY ROSE O'HALLORAN.

(Continuation of observations in No. 71 of these *Publications*.)

- Oct. 22. α *Ceti* is slightly less than ξ *Piscium* in brightness.
 28. Ditto.
 31. Scarcely brighter than ρ *Ceti* on a clear dark sky.
 Nov. 21. Equal to γ *Ceti*.

- 29. It seems less than 7r.
 - 30. It seems equal to 7r.
 - Dec. 1. Less than 7r.
 - 2. Ditto.
 - 5. Ditto.
 - 8. Equal to the star of 8.2, magnitude south of 7r.
 - 17. It is less than 8.2, but brighter in moonlight than adjacent star of 8.4 magnitude.
-
- Oct. 22. *W Lyræ* is brighter than an adjacent star classed as of 12th magnitude, and equal to one classed as 10th.
 - 23. Ditto.
 - 28. Brighter than 10th magnitude, but not as bright as one to the north classed as 8.6; it is probably of about 9th magnitude.
 - 31. It has undergone a rapid decrease, and is again of 10th magnitude.
 - Nov. 1. Ditto.
 - 4. It is now of 9th magnitude.
 - 6. Ditto.
 - 21. It is brighter than 9th magnitude, but not equal to star of 8.6 magnitude.
 - 22. There is a slight decrease.
 - 29. Probably of 9th magnitude.
 - 30. Ditto.
 - Dec. 1. Ditto.
 - 2. Ditto.
 - 3. Ditto.
 - 5. Ditto.
 - 6. Ditto.
 - 17. It equals a comparison star of 8.6 magnitude.